

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

45 88115

STA/STA

S21
A75 U64
C44

An Analysis of Pear Psylla Populations 1977-79

A free copy of this publication is available from the
Yakima Agricultural Research Laboratory, 3706 West
Nob Hill Boulevard, Yakima, WA 98902.

International Standard Serial Number (ISSN) 0913-3760

Science and Education Administration, Agricultural Reviews and Manuals,
Western Series, No. 24, May 1981

Published by Agricultural Research (Western Region), Science and Education
Administration, U.S. Department of Agriculture, Oakland, Calif. 94612

ABSTRACT

The pear psylla, *Psylla pyricola* Foerst., may be controlled in unsprayed orchards by midseason by the hemipteran predators, *Anthocoris* spp. and *Deraeocoris* spp. In commercial orchards, the impact of predators is minimized because of pesticide usage. Application of pesticides for pear psylla and other insects and mites decimates populations of all pear psylla parasites and predators. In some instances, single applications of synthetic pyrethroid insecticides, directed at the overwintering overwintered pear psylla females, successfully controlled pear psylla through harvesttime and allowed reentrance of predators in late season. Extended warm weather during the falls of 1978 and 1979 enhanced late-season buildup of large overwintering populations of pear psylla that were not controlled by attending predator populations.

KEYWORDS: Pear psylla, *Psylla pyricola*, insecticides, pears, pear pest management, predators, *Anthocoris*, *Deraeocoris*, *Phytocoris*, *Nabis*, *Campylomma*, *Chrysopa*, *Hemirrobis*, ladybird beetles, developmental suppression, high temperatures.

This paper contains the results of research only. Mention of pesticides does not constitute a recommendation for use, nor does it imply that the pesticides are registered under the Federal Insecticide, Fungicide, and Rodenticide Act as amended. The use of trade names in this publication does not constitute a guarantee, warranty, or endorsement of the products by the U.S. Department of Agriculture.

CONTENTS

| | Page |
|---|------|
| Introduction..... | 1 |
| Methods..... | 1 |
| Results..... | 2 |
| * Discussion..... | 5 |
| Literature cited..... | 7 |
| Appendix..... | 9 |
| Pesticide designations..... | 9 |
| Orchard descriptions, horticultural practices, insecticide applications, psylla densities, and beating tray results (bar graphs)..... | 10 |

AN ANALYSIS OF PEAR PSYLLA POPULATIONS, 1977-79¹

By R. E. Fye²

INTRODUCTION

The low return on large production investments in the late 1970's has forced many pear growers to review the various facets associated with production. Their analyses have shown that costs of controlling pear insect pests (as much as \$250 per acre) severely reduced the net return. The major pear insect pest requiring control was the pear psylla, *Psylla pyricola* Foerst. The pear psylla problem in California, the Pacific Northwest, and British Columbia, Canada, has been discussed by several authors in the past two decades (1, 2, 7, 21).³

The feeding of the nymphs of the pear psylla results in much honeydew, which drips on the pear fruit, becomes overgrown with sooty mold, and causes a severe russetting that reduces the quality of the pears. In large sustained infestations, the pear trees may be debilitated with a resulting loss of vigor and yield. This study was conducted to determine why certain growers had highly successful control programs for the pear psylla while others had marginal or ineffective controls. The grower programs were highly variable in approach and results because dissimilar climatic, soil, and cultural conditions exist in the various parts of the Yakima Valley.

METHODS

A preliminary evaluation of the pear psylla and the attendant predators populations was made in 1977. In 1978 and 1979, the evaluations were continued in more detail. In 1978, five commercial orchards were monitored to determine the insect populations on the trees and population responses to grower management. In 1979, the insect populations in seven commercial orchards, including the five monitored in 1978, were assessed.

¹ Research financed in part by a grant from the Washington State Tree Fruit Research Commission.

² Research entomologist, Yakima Agricultural Research Laboratory, Yakima, Wash. 98902.

³ Italic numbers in parentheses refer to Literature Cited, p. 7.

The insect populations were sampled at weekly intervals from early March through October or until the foliage had senesced so that it could no longer support the pear psylla. Before the trees had leafed out, 25 fruit spurs in each orchard were examined for the eggs of pear psylla. When the leaves had opened adequately, three leaves from vegetative spurs and three from fruit spurs were examined from each of 25 trees per orchard. When new shoot growth commenced, three young leaves, taken randomly along the shoot, were also examined. The immature psylla populations were determined by visual examination with a microscope. In addition, the populations of pear psylla adults and predators in pear trees were evaluated by the beating tray method (3). Fifteen trays (three limb taps over each 45- x 45-cm tray) were taken in each orchard each week. Therefore, the total sample within each orchard each week consisted of 225 leaves to determine the immature psylla population and 15 beating trays to determine the adult pear psylla and predator populations. At the end of the season, when the foliage became mature and the leaves dropped, the leaf samples were discontinued, but the beating tray samples were continued into November.

RESULTS

Descriptions of the orchards, notes on the horticultural practices, and the insecticide programs of the various growers are presented in the Appendix. The immature psylla densities and beating tray results are presented in bar graphs utilizing a slightly modified logarithmic scale to better delineate the variations in the populations at low levels.

The predators include the larval, nymphal, and adult stages of *Anthocoris* spp., *Deraeocoris* spp., *Campylomma verbasci* (Meyer), *Nabis* spp., *Phytocoris* spp., *Chrysopa* spp., *Hemerobius* spp., earwigs, *Coccinella transversoguttata* Fald., *Adalia bipunctata* (Linn.), and spiders.

The 1977 data from the laboratory orchard (fig. 1)⁴ show a large egg deposition by the overwintering females. In late April and early May, however, an apparent influx of overwintering predators, mainly *Anthocoris melanocerus* Reuter, maintained the immature pear psylla population at less than one per leaf until early June and then departed. The apparent maintenance of low pear psylla levels during late July and early August may be attributed to extended periods of relatively hot, dry weather (13, 14).

In 1978, a late emergence or buildup of predators resulted in the survival of a large number of the first generation of pear psylla (fig. 2). As a consequence, the adult psylla population remained relatively high throughout the season, and the early, heavier, immature population precluded midseason control by the limited number of predators. When the population of immature pear psylla was reduced to about one per leaf in late July, however, the predators departed, and the fall population of adults attained high levels due to the minimal predation on the immatures.

⁴All figures are shown in the Appendix.

In 1979, the large spring egg deposition and resulting nymphs were controlled early in the season by a large early emerging population of predators, mainly *A. melanocerus* (fig. 3). Again, the population of immature pear psylla was reduced to a level that resulted in the departure of the predators, but this occurred after the immature population had been reduced considerably lower than it was in 1978. Increased survival of a reduced number of immature pear psylla, due to relaxed predator pressures, resulted in a buildup of overwintering adults in the fall. In both 1978 and 1979, an extended warm fall enabled a large number of the pear psylla to mature.

The data for the 3 years indicate that the naturally occurring predators, if present in adequate numbers, will effectively control the pear psylla in midseason. Unfortunately, due to excessive fruit drop caused by a large population of codling moths, *Laspeyresia pomonella* (Linn.), we have been unable to determine if the pear psylla was controlled before damage occurred to the fruit. The late appearance of predators in 1978 also indicates that a late emergence and slow buildup of the predators will permit a large population of pear psylla to develop before the predators become effective.

The relatively heavy insecticide program of grower 1 (figs. 4 and 5) demonstrates the effectiveness of the two major insecticides currently used to control the pear psylla. Two early applications of fenvalerate⁵ with two subsequent midsummer applications of amitraz controlled the pear psylla until harvest. When insecticidal pressure was relaxed, the survivors of the summer applications of insecticide were the genesis of a large fall population that overwintered. The absence of predator pressure, due to the lack of a major reservoir in an adjacent unsprayed area, may be a contributing factor to the major fall buildup. The need to apply additional miticides to control destructive mites indicates the suppression of predatory mites by the pesticides.

For more than 5 years, grower 2 had employed minimal applications of insecticide directed toward the pear psylla. In 1978, a similar program maintained the pear psylla at a low level but permitted a late fall buildup of overwintering adults (fig. 6). Although predators, mainly spiders, were present in relatively large populations in late season they failed to control the fall population increase. Likewise, the general absence of predators in the early season indicates that predators were not a major control factor. The control of a minimal overwintering population of adult pear psylla with Q-137 (see p. 10) was probably responsible for the slow-building population.

In 1979, the heavier overwintering population (fig. 7) was not as readily controlled with insecticides, and a stronger population survived. The surviving pear psylla coupled with an increasing population of codling moths and red mites necessitated a midseason application of pesticides that resulted in a severe reduction in the slowly developing population of predators, thus minimizing their impact in the late season. Consequently, the fall population of overwintering individuals was about double that of 1978. The 1978 data clearly demonstrate the benefits of the use of pesticides that are not broad spectrum insecticides;

⁵See section on Pesticide Designations in the Appendix for trade name and chemical name.

however, the 1979 data show the potential rapid regrowth of populations of pear psylla after summer applications of insecticides.

Grower 3 employed a single dormant spray of fenvalerate that resulted in economic control for the entire season (fig. 8). There was a gradual buildup of the immature pear psylla, resulting in a population of adults that was of some concern when harvest approached. The damage from honeydew was apparently reduced and minimized by the overhead sprinkler system (20). The entry of a relatively large population of *Deraeocoris brevis* in late July suggested that an undetected major reservoir of the predator existed in the area. Subsequently, the population of *Deraeocoris* reduced the immature population in late August, September, and early October. The departure of the *Deraeocoris* population to hibernation sites, or in response to the lack of adequate prey, resulted in some survival of the immatures and a slight buildup of adults in late fall; however, the populations were smaller than those in orchards employing intense insecticide control programs (figs. 5, 10, and 12). The data indicate the benefits of a minimal insecticide program, the migratory ability of major predator, and the strong reproductive potential of those predators, particularly *D. brevis*.

In 1978, the control programs of grower 4 showed the effectiveness of strong early season control of pear psylla (fig. 9); however, in late July an application of amitraz was required. In the absence of additional applications, the survivors initiated large late-season populations. Predator pressures in late season were low due to the lack of a major reservoir (in an adjacent unsprayed area) to furnish the initiating populations. In 1979, the early populations of immatures were reduced by insecticide applications (fig. 10), but the survivors of the summer insecticide applications initiated a large potential population of overwintering adults in late fall. The predators were decimated by the insecticide applications and never recovered. The data demonstrate the rapid recovery of pear psylla populations when small numbers survive insecticide applications. The necessity for miticide applications indicated some weakness in the early season control or a reservoir of mites for reinfestation of the trees.

The 1978 early season control program of grower 5 controlled the overwintering adults and early season immatures of the pear psylla (fig. 11). Two applications of a phosmet-ethion combination permitted a slow buildup of the pear psylla in the midseason and resulted in very slight damage to the harvested fruit. After harvest, there was a rapid buildup with a resulting large overwintering population. Predation was ineffective due to the lack of predators in early season and the absence of an unsprayed predator reservoir. The results in 1979 were similar to those in 1978, but the late-season buildup of overwintering forms was slightly less (fig. 12). The general overwintering populations in this orchard were larger than those with fewer applications of insecticides.

An early application of fenvalerate reduced the population overwintering adults and their progeny in the orchard of grower 6 (fig. 13). The low levels were maintained throughout midseason with July applications of detergent employed to minimize honeydew damage. Predator populations were generally low until late July and declined fairly rapidly in late September, resulting in a slight increase in the overwintering population of adults. The population of overwintering adults was lower than that in the orchards employing heavy pesti-

cide programs for the control of pear psylla (figs. 5, 10, and 12). A mite problem that arose in June was effectively controlled with cyhexatin.

In 1978, grower 7 employed a relatively intensive insecticide program that failed to control the pear psylla in midseason (fig. 14). A late-season application of phosmet had relatively little effect on the pear psylla population. After harvest, the pear psylla population increased rapidly, resulting in a large population of overwintering adults. The insecticide applications minimized the predator population in early season, but some recovery occurred from the latter part of July into October. The data emphasized the reproductive potential of the pear psylla and the capability of the insect for rapid recovery to high levels of infestation.

The data presented in figure 15 represent the populations of pear psylla and predators in pear trees interspersed in an apple block. The cover spray program for apples that the trees received aggravated the pear psylla population, and a large population infested the trees from late June through the remainder of the season. Any fruit borne on these trees would have been severely damaged. In late season, there was a strong buildup of predators, mainly *A. melanocerus*. The trees were removed at the conclusion of the 1978 season.

In 1979, the large population of overwintering pear psylla in the orchard of grower 7 was not fully controlled by the early season applications of insecticide (fig. 16). Subsequently, the summer applications left a number of survivors, which resulted in a large late-summer and overwintering population in the fall. The predator populations were controlled by insecticides throughout the summer; however, they recovered slightly in the fall when applications ceased. Again, a major overwintering population was produced in an orchard with a heavy summer insecticide history. The data also demonstrate the inability of late-season predation to make inroads into large populations of potentially overwintering pear psylla.

DISCUSSION

The 2-year study of pear orchards provides some insight into the results of current pear psylla control programs and documents a number of generally accepted ideas. Native *Anthocoris* and *Deraeocoris* have long been known to be effective predators of the pear psylla (4, 6, 8, 9, 10, 11, 12, 15, 17, 18, 19, 22, 22, 23, 24). In the current studies, *Anthocoris* spp. were the major predators during the midseason, and *Deraeocoris* was generally the major predator after the pears had been harvested.

Although the predators provided midseason control in unsprayed situations, they failed--through late emergence or slow buildup--to control the early immature forms of the pear psylla. In addition, their assumed propensity to leave an area when food supplies become inadequate permitted large fall buildups of overwintering pear psylla adults; however, late-season populations of predators, particularly *Deraeocoris*, have the capability of expanding greatly if an unsprayed reservoir is available to provide an initiating population after the insecticide applications are terminated. Other biocontrol organisms may reach population levels that provide some assistance to the major predators. These in-

clude *Trechnites insidiosus*, *Chrysopa* spp., *Adalia bipunctata*, and the spider complex. In localized situations, these parasites and predators have considerable impact upon the pear psylla (24).

High temperatures may have some impact on pear psylla population (13, 14). During the 1978 and 1979 season, the following periods of high temperatures occurred in the Yakima Valley:

| | Number of hours with temperatures over: | | | | | |
|--------|---|------|--------|------|------|------|
| | 30°C | | 32.5°C | | 35°C | |
| | 1978 | 1979 | 1978 | 1979 | 1978 | 1979 |
| April | 7.0 | 0 | 3.5 | 0 | 0 | 0 |
| May | 0 | 0 | 0 | 0 | 0 | 0 |
| June | 49.0 | 23.5 | 11.0 | 2.5 | 0 | 0 |
| July | 58.4 | 58.0 | 17.5 | 19.5 | 1.5 | 0 |
| August | 147.5 | 9.5 | 82.5 | 0 | 19.5 | 0 |

Although some heat suppression of the pear psylla may have occurred during August of 1978, it was probably not extensive. In 1979, however, there was little indication that the high temperatures and not predation were responsible for the maintenance of relatively low levels of pear psylla (fig. 3). Additional research must be conducted to determine the full impact of heat suppression on pear insect populations.

The results from the grower insecticide programs indicate that the two materials currently available, fenvalerate and amitraz, are effective against the early season and midseason pear psylla populations, respectively. The need for thorough application is evident in the rapid rebuilding of the pear psylla populations to high levels after insecticide applications are terminated. A small number of survivors is adequate to initiate a large overwintering population. Hopefully, the low level survival in midseason (figs. 4, 5, 9, and 10) is due to escape from poor insecticide application and migration and not due to the pear psylla's propensity for rapid development of resistance to pesticides.

The recovery of populations of predators after single thorough applications of fenvalerate in the spring suggests a strong potential for integrated control; however, the potential is nullified when applications of pesticides are made for other pests, particularly codling moth and mites. An unsprayed reservoir is essential for providing a source for reestablishing predator populations.

The large population increases in the pear psylla after the harvest of the fruit, when pesticide applications are discontinued, presents a paradox. A fall insecticide program for control of the overwintering population was suggested early (5) and is suggested in the current spray guide for eastern Washington (16).

The success of such programs may be seriously jeopardized by the migratory tendencies of the pear psylla and therefore requires that a large area be treated. Generally, the grower with an intense pesticide program is faced with a large fall population and is placed in the unresolved dilemma of applying pesticides to the overwintering generation in the fall, thus increasing the hazard of the potentially insecticide resistant pear psylla while still facing reinvasion of his orchard by migratory pear psylla in the spring.

The impact of overhead sprinkling systems on pear psylla has been investigated by Westigard et al. (20). They concluded that a major effect of overhead sprinklers is to reduce the damage by the pear psylla honeydew. Although detailed data on the impact of sprinklers were not included in our study, the tendency for minimized russetting damage in the two orchards with overhead sprinklers (figs. 6, 7, and 8) substantiates their observations.

The practice of interplanting pears with apples has generally been discontinued, but the data presented in figure 16 confirmed the need for separate programs for the two fruit. Obviously, pear psylla control in interplanted apples and pears requires adequate separation of trees for selective spraying.

The study suggests that judicious use of the available insecticides remains the strongest base for a prudent pear psylla control program; however, a program employing a number of practices, involving minimal use of hard pesticides, may become feasible if the available materials are lost to the grower for practical or regulatory reasons.

LITERATURE CITED

- (1) Burts, E. C.
1959. Insect control problems on pears in central Washington. Proceedings of the Washington State Horticultural Association 55:161-164.
- (2) _____
1970. Pear psylla in central Washington. Washington Agricultural Experiment Station, Circular No. 516, 13 p.
- (3) _____ and A. H. Retan.
1973. Detection of pear psylla. Washington State University Cooperative Extension Service, Extension Manual No. 3069, 2 p.
- (4) Fields, G. J., and B. P. Beirne.
1973. Ecology of Anthocorid (Hemiptera:Anthocoridae) predators of the pear psylla (Homoptera:Psyllidae) in Okanagan Valley, British-Columbia, Canada. Journal of the Entomological Society of British Columbia 70: 18-19.
- (5) Hodgkiss, H. E.
1913. Fall spraying for pear psylla. Journal of Economic Entomology 6(2):243-244.

- (6) Madsen, H. F.
1961. Notes on *Anthocoris melanocerus* Reuter (Hemiptera:Anthocoridae) as a predator of the pear psylla in British Columbia. Canadian Entomologist 93:660-662.
- (7) _____ R. L. Sisson, and R. S. Bethell.
1962. The pear psylla in California. California Agricultural Experiment Station Circular No. 510, 11 p.
- (8) _____ P. H. Westigard, and R. L. Sisson.
1963. Observations on the natural control of the pear psylla, *Psylla pyricola* Foerster, in California. Canadian Entomologist 95(8):837-844.
- (9) _____ and T. T. Y. Wong.
1964. Effects of predators on control of pear psylla. California Agriculture 18(2):2-3.
- (10) McMullen, R. D.
1964. Biological control of pear psylla. Proceedings of the Washington State Horticultural Association 60:79-82.
- (11) _____ and C. Jong.
1967. The influence of three insecticides on predation of the pear psylla, *Psylla pyricola*. Canadian Entomologist 99:1292-1297.
- (12) _____
1967. New records and discussion of predators of the pear psylla, *Psylla pyricola* Foerster, in British Columbia. Journal of the Entomological Society of British Columbia 64:35-40.
- (13) _____
1972. Influence of temperature and host vigor on fecundity of the pear psylla (Homoptera:Psyllidae). Canadian Entomologist 104:1209-1212.
- (14) _____
1977. Effect of temperature on developmental rate and fecundity of the pear psylla, *Psylla pyricola* (Homoptera:Psyllidae). Canadian Entomologist 109(2):165-169.
- (15) Nickel, J. L., J. T. Shimizu, and T. T. Y. Wong.
1965. Studies on natural control of pear psylla in California. Journal of Economic Entomology 58(5):970-976.
- (16) Washington State University Cooperative Extension Service.
1980. Spray guide for tree fruit in eastern Washington. Extension Bulletin No. 0419, 56 p.
- (17) Watson, T. K., and W. H. A. Wilde.
1963. Laboratory and field observations on two predators of the pear psylla in British Columbia. Canadian Entomologist 95:435.

- (18) Westigard, P. H.
1973. The biology of and effect of pesticides on *Deraeocoris brevis piceatus* (Heteroptera:Miridae). Canadian Entomologist 105:1105-1111.
- (19) _____ L. G. Gentner, and D. W. Berry.
1968. Present status of biological control of the pear psylla in southern Oregon. Journal of Economic Entomology 61(3):740-743.
- (20) _____ P. B. Lombard, and R. B. Allen.
1979. Effects of overtree irrigation on density and damage of pear pests. Journal of Economic Entomology 72:839-840.
- (21) _____ and R. W. Zwick.
1972. The pear psylla in Oregon. Oregon Agricultural Experiment Station Technical Bulletin No. 122, 22 p.
- (22) Wilde, W. H. A.
1962. Bionomics of the pear psylla, *Psylla pyricola* Foerster, in pear orchards of the Kootenay Valley of British Columbia. Canadian Entomologist 94:845-849.
- (23) _____ and T. K. Watson.
1963. Bionomics of the pear psylla, *Psylla pyricola* Foerster, in the Okanagan Valley of British Columbia. Canadian Journal of Zoology 41(6):953-961.
- (24) Zwick, R. W., and G. E. J. Fields.
1977. Integrated control of pear psylla in Oregon's Hood River Valley. Oregon Agricultural Experiment Station, Circular of Information No. 660, 8 p.

APPENDIX

Pesticide Designations

| Trade name | Common name | Chemical name |
|-----------------|---------------|--|
| BAAM® | amitraz | (<i>N,N'</i> -[(methylimino)dimethylidyne] bis[2,4-xylidine]). |
| Diazinon® | diazinon | <i>O,O</i> -diethyl <i>O</i> -(2-isopropyl-6-methyl-4-pyrimidinyl)phosphorothioate. |
| Ethion | ethion | <i>O,O,O',O'</i> -tetraethyl <i>S,S'</i> methylene bis-(phosphorodithioate). |
| Ethyl parathion | parathion | <i>O,O</i> -diethyl <i>O</i> -(<i>p</i> -nitrophenyl)phosphorothioate. |
| Guthion® | azinhosmethyl | <i>O,O</i> -dimethyl <i>S</i> -[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate. |

| <i>Trade name</i> | <i>Common name</i> | <i>Chemical name</i> |
|-------------------|---------------------|---|
| Imidan® | phosmet | <i>O,O</i> -dimethyl phosphorodithioate <i>S</i> -ester with <i>N</i> -(mercaptomethyl)phthalimide. |
| Karathane® | dinocap | 2-(1-methylheptyl)-4,6-dinitrophenyl crotonate (commercial product is a reaction mixture containing several isomers). |
| Lime sulfur | calcium polysulfide | 30-percent calcium polysulfide and various small amounts of calcium thiosulfate plus water and free sulfur. |
| Morestan® | oxythioquinox | cyclic <i>S,S</i> -(6-methyl-2,3-quinoxalinediyl) dithiocarbonate. |
| Perthane® | Q-137 | 1,1-dichloro-2,2-bis(<i>p</i> -ethylphenyl)ethane. |
| Plictran® | cyhexatin | tricyclohexylhydroxystannane. |
| Pydrin® | fenvalerate | cyano(3-phenoxyphenyl)methyl 4-chloro- α -(1-methyl=ethyl)benzeneacetate. |
| Solabar® | barium polysulfide | |
| Superior oil | petroleum oil | |
| Thiodan® | endosulfan | 6,7,8,9,10,10-hexachloro-1,5,5a,6,9,9a-hexahydro-6,9-methano-2,4,3-benzodioxathiepin 3-oxide. |

Orchard Descriptions, Horticultural Practices, Insecticide Applications, Psylla Densities, and Beating Tray Results (Bar Graphs)

Yakima Agricultural Research Laboratory Orchard (figs. 1-3).

Location: Yakima.

Situation: Mature orchard (1 acre) with scattered replacement trees of variable age. Dwellings on south and west; laboratory on north, vegetable experimental plots on east beyond 2 acres of apples.

Irrigation: Rill.

Ground cover:

1977 - Light sod with strong admixture of weeds.




Figure 1.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of predators per tray. Yakima Agricultural Research Laboratory, Yakima, Wash., 1977.




Figure 2.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of adult pear psylla and predators per tray. Yakima Agricultural Research Laboratory, Yakima, Wash., 1978.




Figure 3.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of adult pear psylla and predators per tray. Yakima Agricultural Research Laboratory, Yakima, Wash., 1979.

Ground cover: (Continued)

1978 - Early light sod with strong admixture of weeds. After July 15, disked, mostly bare soil. After August 15, experimental plantings of wheat and barley planted at weekly intervals until September 5.

1979 - Fall and spring planted wheat with heavy admixture of weeds. Grass and weeds in the tree root area.

Insecticide applications:

1977-79 None

Grower 1 (figs. 4 and 5)

Location: Yakima. North slope of Ahtanum Ridge.

Situation: Bearing orchard (about 50 acres) of even aged trees. Open field on north, bearing pear orchard on east, open field on south, open field on west. Some rows of d'Anjou pollinators.

Irrigation: Rill.

Ground cover: Cultivated, with small amounts of weeds escaping herbicide applications.

Insecticide applications:

1978

| | | |
|----------|-------------------------|------------------------------|
| March 1 | Superior Oil Pydrin® | 4 gal/acre 0.3 lb AI/acre |
| March 24 | Pydrin® | 0.25 lb AI/acre |
| May 23 | Guthion® | 0.9 lb AI/acre |
| July 10 | BAAM® | 1.1 lb AI/acre |
| | Plictran® | 1.0 AI/acre |
| | Guthion® | 0.6 AI/acre |
| August 1 | BAAM® | 0.8 lb AI/acre |
| | Guthion® | 0.75 AI/acre |

1979

| | | |
|----------|-------------------------|------------------------------|
| March 13 | Superior Oil Pydrin® | 5 gal/acre 0.2 lb AI/acre |
| April 6 | Pydrin® | 0.2 lb AI/acre |
| May 2 | Plictran® | 1.0 lb AI/acre |
| June 25 | BAAM® | 1.1 lb AI/acre |
| | Plictran® | 0.75 lb AI/acre |




Figure 4.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of adult pear psylla and predators per tray. Grower 1, Yakima, Wash., 1978.




Figure 5.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of adult pear psylla and predators per tray. Grower 1, Yakima, Wash., 1979.

Insecticide applications: (Continued)

1979

| | | |
|---------|-----------|-----------------|
| July 27 | BAAM® | 0.8 lb AI/acre |
| | Plictran® | 0.75 lb AI/acre |
| | Guthion® | 0.75 lb AI/acre |

Grower 2 (figs. 6 and 7)

Location: 4 miles north of Zillah.

Situation: Mixed orchard of bearing Bartlett and d'Anjous (5 acres). Row crops and alfalfa on east. Planted to apples in 1979. Five acres of apples on south. Asparagus on west, open range on north. Slight slope facing west.

Irrigation: Overhead sprinkler.

Ground cover: Dense weeds.

Insecticide applications:

1978

| | | |
|------------|--------------|----------------|
| March 22 | Superior Oil | 1 gal/acre |
| | Perthane® | 4 lb AI/acre |
| | Parathion | 1.5 lb AI/acre |
| May 24 | Diazinon® | 2.5 lb AI/acre |
| June 20 | Diazinon® | 2.5 lb AI/acre |
| July 19 | Diazinon® | 2.5 lb AI/acre |
| October 19 | Lime-sulfur | 6 gal/acre |

1979

| | | |
|-----------------------------|--------------|-----------------|
| March 17 | Superior Oil | 5 gal/acre |
| | Perthane® | 2 lb AI/acre |
| March 23 | Superior Oil | 8 gal/acre |
| | Parathion | 0.75 AI/acre |
| | Thiodan® | 2.25 AI/acre |
| April 28 (d'Anjous only) | Karathane® | 0.3 lb AI/acre |
| May 20 | Guthion® | 1.25 lb AI/acre |
| July 19 | BAAM® | 0.60 lb AI/acre |
| | Parathion | 0.75 lb AI/acre |
| | Plictran® | 0.75 lb AI/acre |




Figure 6.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of adult pear psylla and predators per tray. Grower 2, Zillah, Wash., 1978.




Figure 7.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of adult pear psylla and predators per tray. Grower 2, Zillah, Wash., 1979.

Grower 3 (fig. 8)

Location: 4 miles north of Zillah.

Situation: Young orchard nearing maturity with interplanted younger trees. Open range land on east, mature apple orchards on north and south, newly prepared orchard land on west. Strong slope facing west.

Irrigation: Overhead sprinkler.

Ground cover: Quack grass with strong admixture of dandelions and other broadleafed weeds.

Insecticide applications:

1979

| | | |
|-------------|-------------------------|------------------------------------|
| March 13-14 | Pydrin® Superior Oil | 0.2 lb AI/acre 3 gal/acre |
| June 1 | Guthion® Plictran® | 0.63 lb AI/acre 1.00 lb AI/acre |

Grower 4 (figs. 9 and 10)

Location: West Wapato area in Lower Yakima Valley. Near Coe and Lateral 1.

Situation: Bearing orchard (4.5 acres) of Bartlettts with d'Anjou pollinators. Apples on north, south, and west. Wasteland east.

Irrigation: Rill.

Ground cover: Cultivated early with increasing cover of weeds as season progressed.

Insecticide applications:

1978

| | | |
|----------|--|---|
| March 7 | Superior Oil Perthane® | 4 gal/acre 4 lb AI/acre |
| March 26 | Parathion Superior Oil Lime sulfur | 0.5 AI/acre 3 gal/acre 3 gal/acre |
| July 11 | Plictran® | 0.75 lb AI/acre |
| July 20 | BAAM® Guthion® | 1.1 lb AI/acre 1.0 AI/acre |




Figure 8.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of adult pear psylla and predators per tray. Grower 3, Zillah, Wash., 1979.




Figure 9.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of adult pear psylla and predators per tray. Grower 4, Wapato, Wash., 1978.




Figure 10.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of adult pear psylla and predators per tray. Grower 4, Wapato, Wash., 1979.

Insecticide applications: (Continued)

1979

| | | |
|----------|-------------------------|------------------------------------|
| March 12 | Pvdrin® Superior Oil | 0.3 lb AI/acre 4 gal/acre |
| April 7 | Morestan® Parathion | 0.63 lb AI/acre 0.75 lb AI/acre |
| June 15 | Plictran® Imidan® | 0.5 lb AI/acre 1.5 lb AI/acre |
| July 18 | BAAM® Imidan® | 0.3 lb AI/acre 1.5 lb AI/acre |

Grower 5 (figs. 11 and 12)

Location: Cowiche.

Situation: Triangular orchard (7 acres) with large cold storage facility on northeast, open area with dwellings and farm buildings on south, apple orchard on west. Bartlett trees of uneven age from young trees to mature. Some d'Anjous and trees from other cultivars.

Irrigation: Rill.

Ground cover: Clean cultivation early with increasing weed stand to harvesttime.

Insecticide applications:

1978

| | | |
|----------|-------------|----------------|
| March 22 | Pydrin® | 0.3 lb AI/acre |
| April 14 | Lime-sulfur | 12 gal/acre |
| July 6 | Imidan® | 3 lb AI/acre |
| | Ethion | 1.5 lb AI/acre |
| July 24 | Imidan® | 3 lb AI/acre |
| | Ethion | 1.5 lb AI/acre |

1979

| | | |
|----------|-----------------------|--------------------------------|
| March 21 | Pydrin® | 0.3 lb AI/acre |
| April 25 | Morestan® Solabar® | 1 lb AI/acre 4 lb AI/acre |
| June 20 | Imidan® Ethion | 3 lb AI/acre 1.5 lb AI/acre |




Figure 11.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of adult pear psylla and predators per tray. Grower 5, Cowiche, Wash., 1978.




Figure 12.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of adult pear psylla and predators per tray. Grower 5, Cowiche, Wash., 1979.

Insecticide applications: (Continued)

1979

| | | |
|---------|-----------|----------------|
| July 19 | Imidan® | 3 lb AI/acre |
| | Ethion | 1.5 lb AI/acre |
| | Plictran® | 0.5 lb AI/acre |

Grower 6 (fig. 13)

Location: Tieton.

Situation: Orchard (4.0 acres) about 15 years old with interplanted young trees. Open rangeland on south and west, mature apples orchard on north, with young and mature apple orchard on east. Weedy ground cover, mainly dandelions and field bindweed, early in season. Heavily rototilled in midsummer with considerable regrowth in late summer and fall.

Irrigation: Undertree sprinkler.

Insecticide applications:

1979

| | | |
|---------|------------------------------|----------------|
| March 7 | Superior Oil | 2 gal/acre |
| | Pydrin® | 0.3 lb AI/acre |
| June 26 | Plictran® | 1.0 lb AI/acre |
| July 3 | Detergent (White King D®) | 10 lb/acre |
| July 6 | Detergent (White King D®) | 10 lb/acre |
| July 27 | Detergent (White King D®) | 10 lb/acre |

Grower 7a and b (figs. 14-16)

Location: Naches.

Situation:

- (a) Ten acres of uneven age trees from recent replacements to scattered mature trees. North end abuts railroad right-of-way with rangeland beyond. Young apple orchard to east. Mature apple orchard on south with bearing peach orchard on west.
- (b) Scattered young bearing pear trees in bearing apple orchard.

Irrigation: Ground level hand lines--sprinkler.




Figure 14.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of adult pear psylla and predators per tray. Grower 7a, Naches, Wash., 1978.




Figure 15.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of adult pear psylla and predators per tray in pears interplanted with apples. Grower 7b, Naches, Wash., 1978.




Figure 16.--Numbers of pear psylla eggs and nymphs per fruiting spur (early season) or leaf (late season) and numbers of adult pear psylla and predators per tray. Grower 7, Naches, Wash., 1979.

Ground cover:

- (a) Thick sod with strong complement of dandelions.
- (b) Thick sod intermixed with red clover.

Insecticide applications:

1978a

| | | |
|----------|--------------|-----------------|
| March 13 | Pydrin® | 0.3 lb AI/acre |
| March 25 | Superior Oil | 2 gal/acre |
| June 5 | Ethion | 1.5 lb AI/acre |
| | BAAM® | 0.75 lb AI/acre |
| July 12 | Ethion | 1.5 lb AI/acre |
| | BAAM® | 0.75 lb AI/acre |
| August 1 | Imidan® | 3 lb AI/acre |

1978b - Cover spray program for apples

| | | |
|----------------|----------|----------------|
| 2 applications | Guthion® | 2.0 lb AI/acre |
|----------------|----------|----------------|

1979

| | | |
|----------|---------------------|-----------------|
| March 17 | Pydrin® | 0.3 lb AI/acre |
| April 5 | Superior Oil | 2 gal/acre |
| | Calcium polysulfide | 10 lb/acre |
| May 25 | Plictran® | 0.75 lb AI/acre |
| June 10 | Imidan® | 2.5 lb AI/acre |
| July 10 | Imidan® | 2.5 lb AI/acre |
| | Ethion | 1.25 lb AI/acre |
| August 2 | Imidan® | 2.5 lb AI/acre |
| | Plictran® | 0.63 lb AI/acre |

* U.S. GOVERNMENT PRINTING OFFICE: 1981-791-212/199

PRINTED MATTER



POSTAGE AND FEES PAID
U. S. DEPARTMENT OF
AGRICULTURE
AGR 101

U. S. DEPARTMENT OF AGRICULTURE
SCIENCE AND EDUCATION ADMINISTRATION
WESTERN REGION
1333 BROADWAY, SUITE 400
OAKLAND, CALIFORNIA 94612
OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300